

61220 – 279 grams
61240 – 452 grams
Immature Trench Soil

DRAFT




Figure 1: Trench dug on rim of Plum Crater, Apollo 16. AS16-109-17801. 61220 was from the white material at the bottom of this trench; 62240 was from the darker material near the top of the trench. Gnomon is 0.5 m.




Figure 2: Proportion diagrams of agglutinates (A), minerals (M) and lithic breccia (L) fragments in various Apollo 16 soils as function of grain size, showing 61221 as unusually low in agglutinates for all size range (from Houck 1982).

Modal content of 61221 and 61241 (90-150 micron).

From Heiken et al. 1973.

	61221	61241
Agglutinates	6.3 %	27.1
Basalt	0.6	2.3
Breccia	33.9	34.5
Anorthosite	13.8	5
Norite	2	-
Gabbro	0.3	0.3
Plagioclase	24.6	19.9
Pyroxene	4.6	4
Olivine	-	-
Ilmenite	-	-
Glass other	12.2	3.3

Introduction

Lunar soil 61220 is different from other lunar soil samples. It is an immature soil that was taken from bottom of a trench dug on the east rim of Plum Crater. The depth is uncertain, but less than 30 cm (figure 1). Soil 61220 is lighter in color than the surface soil (61240) (Morris et al. 1983).

Based on outgassing properties (trace CN) Gibson and Moore (1973) speculated that this soil might include cometary material, but this hypothesis has not been confirmed by other investigators (see, for example, Epstein and Taylor 1973). However, this soil is unusual in several respects and the origin of its volatile content remains an enigma.

Mineralogic Mode for 61221

	LSPET 73	Morris 83	Houck 82
Agglutinates	8	6.3	6.3
Glass fragments			9.9
Colorless	23	10.9	
Brown		1.3	
Olivine	1		0.2
Pyroxene	8	4.6	6.9
Plagioclase	35	17	29.5
Maskeleynite		7.6	
Metabreccia	12	18.3	32.8
Vitric breccia	10	15.6	11.7
Anorthosite		13.6	1.9
Basalt	1	0.6	-
Norite		2	-

Petrography

Karen Houck (1982) determined the detailed mineralogic mode of 61221 and its various grain sizes, finding this sample unusual (figure 2). Houck (1982) and Heiken et al. (1973) found a high proportion of glass, but everyone agrees that the sample is low in agglutinate content. Ridley et al. (1973) reported that it contained a lot of maskelynite (or plagioclase glass). The maturity index, determined by magnetic properties, is low ($Is/FeO = 9.2$) (Morris 1979).

Cadenhead et al. (1977) found that 61221 had a high surface area for its low maturity index, while 61241 was within the normal range (figure 15). Cadenhead et al. speculate that 61221 was created through a single impact event.

Glass

Ridley et al. (1973) and Meyer (1978) studied the glass particles in 61221. Ridley et al. argued that clusters in glass composition allowed them to identify “Highland Basalt” and other “rock types”. However, the

composition of “Highland Basalt” closely matches the composition of the bulk soil (figures 13 and 14). Meyer attempted to show that “average” glass composition is contaminated by trace elements, because at least some of the glass is made from the regolith or from breccia (mixtures).

Coarse Fines

Marvin and Mosie (1980) made a catalog of coarse fines (1-4 mm) from 61220 and these have been subsequently investigated by various labs. Marvin (1972) included coarse-fines from 61240 in her catalog of 4-10 mm particles. Several small rocks from these soils were given individual sample numbers and are briefly described in the Apollo 16 Rock Sample Catalogs by Butler (1972) and Ryder and Norman (1980).

Marvin and Warren (1980) studied the particles of pristine “micro-gabbro” found in the coarse fines (table 3). 61224,6 is composed of ~63% pyroxene (0.2 – 3




Figure 3: Composition diagram for lunar soils showing that 61221 is high aluminous.




Figure 4: Normalized rare-earth-element diagram for 61221 and 61241 (see tables).




Figure 5: 61221 is special in that it has relatively high carbon content and low maturity.

mm) and ~34% plagioclase (0.3 – 1 mm). Plagioclase is An_{83} and pyroxene is $En_{65}Fs_{32}Wo_3$ and $En_{43}Fs_{17}Wo_{40}$. Takeda et al. (1981) studied the nature of the exsolved pyroxene in 61224,6, discussing the cooling history.

Chemistry

The chemical composition of 61221 was originally reported by LSPET (1973), Bansal et al. (1972), Haskin et al. (1973), Rose et al. (1973) and others (table 1). Boynton et al. (1976) provided a complete analysis for 61241 (table 2). Finkelman et al. (1975) and Korotev et al. (1981) also measured the composition of the fine fraction from these samples, and Rhodes et al. (1975) studied the agglutinates in 61241. These soils are highly aluminous (figure 3). 61241 has a higher trace element content than 61221, from the bottom of the trench (tables 1 and 2, figure 4).

Gibson and Moore (1973b) give the sulfur content of 61221 as 496 ppm. Jovanovic and Reed (1973) studied the halogens in 61221 and 61241 finding that they were within the range of other Apollo 16 materials. Allen et al. (1974) found that ^{204}Pb , Bi, Tl and Zn were within the overall range of Apollo 16 soils.

Mark et al. (1973) and Nyquist et al. (1973) determined the Rb, Sr and K contents and Sr isotopic composition for 61221 and 61241.

61221 has relatively high carbon content (~100 ppm) for its maturity (Moore et al. 1973, figure 5). Gibson and Moore (1973a) and Wszolek et al. (1973) studied the temperature release of volatiles from 61221 (figures 6, 7abc). These temperature release patterns are different from that of other lunar materials, with more carbon based molecules coming off at a lower temperature. Gibson and Moore speculated that the volatiles might have a cometary origin, while Wszolek et al. interpreted this unusual volatile pattern might be from an impact by a carbonaceous chondrite. However, DesMarais et al. (1973) found the carbon content and species similar to that of other lunar soils.

Epstein and Taylor (1973) found that the H/D isotope ratio was like that of terrestrial water (figure 10) and provide an indepth discussion of the H, C and oxygen isotope measurements with respect to Gibson and Moore's cometary hypothesis. Sill et al. (1974) have also suggested a cometary origin for gas trapped in




Figure 6: Temperature release curves for volatiles from 61221 (Gibson and Moore 1973). Note the verticle scale is "arbitrary".




Figure 7a: Temperature release curves for 61221 (from Wszolek et al. 1973).




Figure 7b: Temperature release curves for 61221 (from Wszolek et al. 1973).




Figure 7c: Temperature release curves for 61221 (from Wszolek et al. 1973).

78155. But no firm evidence has been established. Becker et al. (1976) determined the nitrogen isotope ratio as a function of release temperature (figure 9).

Krahenbuhl et al. (1973) and Ganapathy et al. (1974) give concentrations for three small particles from 61222.

Cosmogenic isotopes and exposure ages

Walton et al. (1973) determined the cosmic ray exposure age of 61221 and 61241 by ²¹Ne as 250 m.y. and 240 m.y., respectively. Imamura et al. (1974) studied ⁵⁴Mn in both samples finding that they were consistent with the ⁵⁴Mn depth profiles of deep drill samples. The main conclusion is that both sample are saturated with ⁵⁴Mn activities predicted for their depth and must have remain exposed and relatively undisturbed for >5 m.y. (perhaps the age of Plum Crater).

Fireman et al. (1973) reported ³⁷Ar, ³⁹Ar and ³H, while Fields et al. (1973) studied the interaction of the cosmic-ray flux with U.

Other Studies

Taylor and Epstein (1973) studied the isotopic composition of silicon and oxygen removed by partial fluorination and compared results with surface soils.




Figure 8: Carbon and hydrogen in 61221 compared with other lunar soils (from Epstein and Taylor 1973).




Figure 9: Isotopic composition of nitrogen released from 61221 during stepwise heating (Becker et al 1976).

Bogard and Nyquist (1973), Walton et al. (1973) and Nautiyal et al. (1981) determined rare gas contents of 61221 and found that 61221 had considerable excess ^{40}Ar and ^4He .

Fleischer and Hart (1974) and Rao et al. (1979) determined the density of cosmic-ray tracks in 61221 (figure 12).

Processing

Soils 61220 and 61240 were brought back in ALSRC#1, but it did not seal and the samples were exposed to spacecraft (LM and CM) and Pacific Ocean atmospheres. A fairly large portion of 61220 remains unsieved. There is a note in the inventory that 61220 contains “clods”.

Several small rocks were removed – see table.




Figure 10: Hydrogen isotope diagram for 61221 and other soils (Epstein and Taylor 1973).




Figure 11: Carbon isotope composition of 61221 (from Epstein and Taylor 1973).




Figure 12: Tracks in 61221 compared with tracks measured for 69921 (Rao et al. 1979).




Figure 13: Glass composition in 61221 showing clusters for "Highland Basalt" and maskelynite (Ridley et al. 1973). BULK composition for table 1 is added (for good measure).




Figure 14: Al_2O_3 content of glass particles in 61221 compared with 60501 (Ridley et al. 1973). The "cluster" at 28% Al_2O_3 (Highland Basalt) is suspiciously like the composition of the bulk regolith (line).





Figure 15: 61221 has a high surface area for its maturity (Cadenhead et al. 1977).



Small rocks found in 61220 and 61240

61224,6	0.5 g	Micro-gabbro
61225	3.52 g	Crystalline Impact Melt
61226	1.53	Cataclastic Anorthosite
61245	8.25	Fine-grained Impact Melt
61246	6.05	Fine-grained Impact Melt
61247	2.48	Poikilitic Impact Melt
61248	1.71	Fagmental Polymict Breccia
61249	1.17	Basaltic Impact Melt (?)
61255	1.13	Cindery Glass

Table 1. Chemical composition of 61221.

			Bansal 72		61220												
reference		Krahenbuhl73	Wiesmann 74		Gibson 73	Haskin 73	Rose 73		Korotev 81				Finkelman75				
weight					LSPET 73				150-90	<20 microns			1mm-90	30-90	<30		
SiO2 %			45.4	(c)	45.35	(c)	46.1	(h)	45.2	(e)							
TiO2			0.45	(b)	0.49	(c)	1.05		0.51								
Al2O3			28.25	(c)	28.25	(c)	26.4		28.3								
FeO			4.55	(c)	4.55	(c)	5.71		4.61		4.36	4.63	(f)				
MnO					0.06	(c)	0.058		0.06								
MgO			5.02	(c)	5.02	(c)	6		4.92								
CaO			16.21	(c)	16.21	(c)	15.7		16.02								
Na2O			0.5	(b)	0.42	(c)	0.52		0.42		0.549	0.532	(f)				
K2O			0.086	(c)	0.09	(c)	0.132		0.08								
P2O5					0.1	(c)			0.11								
S %					0.06	(c)											
sum																	
Sc ppm						7	(d)	6.9		7.2	7.7	(f)	8	8	10	(g)	
V								20					14	20	24	(g)	
Cr		526	(b)	590	(c)	600	(d)	615		575	760	(f)					
Co						17	(d)	9.2		24.5	14.5	(f)	10	15	16	(g)	
Ni				109	(c)			160		330	210	(f)	150			(g)	
Cu								6.2					5			(g)	
Zn	21.5	(a)				23	(d)	22					18	21		(g)	
Ga						6.2	(d)	4.6					3	3	4	(g)	
Ge ppb	345	(a)															
As																	
Se	206	(a)															
Rb	2	(a)	1.96	(b)		2.3	(d)	1.8					1			(g)	
Sr			188	(b)	182			145					180	160	170	(g)	
Y								26					28	30	35	(g)	
Zr			110	(b)				85					86			(g)	
Nb																	
Mo																	
Ru																	
Rh																	
Pd ppb																	
Ag ppb	11	(a)															
Cd ppb	140	(a)															
In ppb																	
Sn ppb																	
Sb ppb	1.6	(a)															
Te ppb	14.7	(a)															
Cs ppm	0.09	(a)				0.096	(d)										
Ba			96.5	(b)				88					93			(g)	
La						9.1	(d)			7.2	9.9	(f)			17	(g)	
Ce			21.6	(b)		22.7	(d)			18.5	26.5	(f)					
Pr																	
Nd			13.5	(b)		14.5	(d)										
Sm			3.82	(b)		4.35	(d)			3.35	4.5	(f)					
Eu			1.21	(b)		1.28	(d)			1.19	1.18	(f)					
Gd			4.6	(b)		5.3	(d)										
Tb						0.9	(d)			0.78	0.97	(f)					
Dy			5.12	(b)		5.76	(d)										
Ho						1.1	(d)										
Er			3.06	(b)		3.2	(d)										
Tm																	
Yb			2.77	(b)		3.01	(d)	1.8		2.65	3.3	(f)	3	2.8	3.2	(g)	
Lu			0.41	(b)		0.44	(d)			0.38	0.44	(f)					
Hf			3.3	(b)		3	(d)			2.3	3.1	(f)					
Ta										0.4	0.4	(f)					
W ppb																	
Re ppb	0.578	(a)															
Os ppb																	
Ir ppb	6.21	(a)															
Pt ppb																	
Au ppb	4.94	(a)															
Th ppm			1.56	(b)						1.2	1.4	(f)					
U ppm			0.38	(b)													
technique: (a) RNAA, (b) IDMS, (c) XRF, (d) INAA, (e) combined methods, (f) INAA, (g) OES, (h) AA																	

Table 2. Chemical composition of 61241.

reference weight	LSPET73	Boynton76 261 mg	Haskin73	Krahenbuhl73	Rose73	Finkelman73 1mm-90 30-90	<30						
SiO2 %	45.32	(a)	45.6	(d)	45.14	(e)							
TiO2	0.57	(a)	0.77	(b)	0.55	(e)							
Al2O3	27.15	(a)	25.89	(b)	27.35	(e)							
FeO	5.33	(a)	5.27	(b)	5.42	(e)							
MnO	0.07	(a)	0.06	(b)	0.068	(e)							
MgO	5.75	(a)	5.47	(b)	5.71	(e)							
CaO	15.69	(a)	15.8	(b)	15.9	(e)							
Na2O	0.55	(a)	0.47	(b)	0.5	(e)							
K2O	0.1	(a)	0.1	(b)	0.124	(e)							
P2O5	0.13	(a)			0.11	(e)							
S %	0.07	(a)											
sum													
Sc ppm		9.1	(b)	8.8	(b)	6.9	(e)	10	8	10	(f)		
V		18	(b)			18	(e)	15	18	21	(f)		
Cr	720	(a)	760	(b)	607	684	(e)						
Co			22.4	(b)	23.8	14	(e)	200	16	16	(f)		
Ni	220	(a)	276	(c)		240	(e)	>1000	320	340	(f)		
Cu						8.8	(e)	15	>55	>55	(f)		
Zn			25.4	(c)	23	22.5	(b)	27	(e)	7	90	213	(f)
Ga			5.4	(c)	5.1	4	(e)	3	3	4	(f)		
Ge ppb			620	(c)		582							
As													
Se					257								
Rb	2.7	(a)		1.9	2.2	2.7	(e)	2	6	11	(f)		
Sr	175	(a)				160	(e)	180	160	170	(f)		
Y	37	(a)				38	(e)	40	30	40	(f)		
Zr	162	(a)				117	(e)	110	110	135	(f)		
Nb	9.8	(a)											
Mo													
Ru													
Rh													
Pd ppb													
Ag ppb					11								
Cd ppb		124	(c)		119								
In ppb		13.9	(c)										
Sn ppb								<10	<10	62	(f)		
Sb ppb					1.55								
Te ppb					15.6								
Cs ppm				0.121	0.105								
Ba		120	(b)			105	(e)	78	94	112	(f)		
La		12	(b)	11.8				<10	<10	<10	(f)		
Ce		29	(b)	30.5									
Pr													
Nd				20									
Sm		5.2	(b)	5.85									
Eu		1.1	(b)	1.29									
Gd				7.2									
Tb		1.1	(b)	1.16									
Dy		6.1	(b)	7.2									
Ho				1.4									
Er				4.9									
Tm													
Yb		3.7	(b)	4.08		2.4		3.3	2.9	3.7	(f)		
Lu		0.5	(b)	0.59									
Hf		3.3	(b)	4.5									
Ta		0.4	(b)										
W ppb													
Re ppb					0.762	(c)							
Os ppb													
Ir ppb		7.1	(c)		9.64	(c)							
Pt ppb													
Au ppb		5.2	(c)		5.31	(c)							
Th ppm	1.7	(a)	1.8	(b)									
U ppm		0.52	(b)										
technique: (a) XRF, (b) INAA, (c) RNAA, (d) AA, (e) mixed, (f) OES													

technique: (a) XRF, (b) INAA, (c) RNAA, (d) AA, (e) mixed, (f) OES

Table 3. Chemical composition of 61224,6.

<i>reference</i>	Marvin80
<i>weight</i>	
SiO ₂ %	50.7
TiO ₂	0.4
Al ₂ O ₃	13.2
FeO	9.91
MnO	0.16
MgO	12.8
CaO	11.6
Na ₂ O	0.91
K ₂ O	0.017
P ₂ O ₅	
S %	
<i>sum</i>	
Sc ppm	20.8
V	
Cr	1991
Co	23.6
Ni	8.3
Cu	
Zn	4
Ga	
Ge ppb	4.3
As	
Se	
Rb	
Sr	
Y	
Zr	170
Nb	
Mo	
Ru	
Rh	
Pd ppb	
Ag ppb	
Cd ppb	4.1
In ppb	0.6
Sn ppb	
Sb ppb	
Te ppb	
Cs ppm	
Ba	32
La	1.47
Ce	4.3
Pr	
Nd	9
Sm	0.87
Eu	1.43
Gd	
Tb	0.22
Dy	
Ho	
Er	
Tm	
Yb	1.06
Lu	0.16
Hf	0.55
Ta	0.16
W ppb	
Re ppb	0.012
Os ppb	
Ir ppb	0.15
Pt ppb	
Au ppb	0.08
Th ppm	0.19
U ppm	<0.6
<i>technique:</i>	(a) INAA and RNAA.

References

- Allen R.O., Jovanovic S. and Reed G.W. (1974) A study of 204Pb partition in lunar samples using terrestrial and meteoritic analogs. Proc. 5th Lunar Sci. Conf. 1617-1623.
- Bansal B.M., Church S.E., Gast P.W., Hubbard N.J., Rhodes J.M. and Wiesmann H. (1972) Chemical composition of soil from the Apollo 16 and Luna 20 sites. Earth Planet. Sci. Lett. 17, 29-35.
- Becker R.H., Clayton R.N. and Mayeda T.K. (1976) Characterization of lunar nitrogen components. Proc. 7th Lunar Sci. Conf. 441-458.
- Bogard D.D., Nyquist L.E., Hirsch W.C. and Moore D.R. (1973) Trapped solar and cosmogenic Noble-gas abundances in Apollo 15 and Apollo 16 deep drill samples. Earth Planet. Sci. Lett. 21, 52-69.
- Bogard and Nyquist (1973) 40Ar/39Ar variations in Apollo 15 and 16 regolith. Proc. 4th Lunar Sci. Conf. 1975-1985.
- Boynton W.V., Chou C.L., Robinson K.L., Warren P.H. and Wasson J.T. (1976) Lithophiles, siderophiles and volatiles in Apollo 16 soils and rocks. Proc. 7th Lunar Sci. Conf. 727-742.
- Cadenhead D.A., Brown M.G., Rice D.K. and Stetter J.R. (1977) Some surface area and porosity characterizations of lunar soils. Proc. 8th Lunar Sci. Conf. 1291-1303.
- Charette M.P. and Adams J.B. (1975) Agglutinates as indicators of lunar soil maturity: The rare gas evidence at Apollo 16. Proc. 6th Lunar Sci. Conf. 2281-2289.
- Cirlin E.H. and Housley R.M. (1981) Distribution and evolution of Zn, Cd and Pb in Apollo 16 regolith samples and the average U-Pb ages of the parent rocks. Proc. 12th Lunar Planet. Sci. Conf. 529-540.
- DesMarias D.J., Hayes J.M. and Meinshein W.G. (1973) The distribution in lunar soil of carbon released by pyrolysis. Proc. 4th Lunar Sci. Conf. 1543-1558.
- DesMarias D.J., Hayes J.M. and Meinshein W.G. (1974) The distribution in lunar soil of hydrogen released by pyrolysis. Proc. 5th Lunar Sci. Conf. 1811-1822.
- Epstein S. and Taylor H.P. (1973) The isotopic composition and concentration of water, hydrogen, and carbon in some Apollo 15 and 16 soils and in the Apollo 17 orange soil. Proc. 4th Lunar Sci. Conf. 1559-1575.
- Fields P.R., Diamond H., Metta D.N. and Rokop D.J. (1973) Reaction products of lunar uranium and cosmic rays. Proc. 4th Lunar Sci. Conf. 2123-2130.

- Finkelman R.B., Baedeker P.A., Christian R.P., Berman S., Schnepfe M.M. and Rose H.J. (1975) Trace-element chemistry and reducing capacity of grain size fractions from the Apollo 16 regolith. *Proc. 6th Lunar Sci. Conf.* 1385-1398.
- Fireman E.L., D'Amico J. and DeFelice J. (1973) Radioactivities vs. depth in Apollo 16 and 17 soil. *Proc. 4th Lunar Sci. Conf.* 2131-2144.
- Fleischer R.L. and Hart H.R. (1974) Particle track record of Apollo 16 rocks from Plum Crater. *J. Geophys. Res.* 79, 766-768.
- Ganapathy R., Morgan J.W., Higuchi H., Anders E., and Anderson A.T. (1974) Meteoritic and volatile elements in Apollo 16 rocks. *Proc. 5th Lunar Sci. Conf.* 1659-1684.
- Gibson E.K. and Moore G.W. (1973a) Volatile rich lunar soil: Evidence of possible cometary impact. *Science* 179, 69-71.
- Gibson E.K. and Moore G.W. (1973b) Carbon and sulfur distributions and abundances in lunar fines. *Proc. 4th Lunar Sci. Conf.* 1577-1586.
- Gibson E.K. and Andreas F.F. (1978) Nature of the gases released from lunar rocks and soils upon crushing. *Proc. 9th Lunar Planet. Sci. Conf.* 2433-2450.
- Haskin L.A., Helmke P.A., Blanchard D.P., Jacobs J.W. and Telander K. (1973) Major and trace element abundances in samples for the lunar highlands. *Proc. 4th Lunar Sci. Conf.* 1275-1296.
- Heiken G.H., McKay D.S. and Fruland R.M. (1973) Apollo 16 soils – grain size analysis and petrography. *Proc. 4th Lunar Sci. Conf.* 251-266.
- Heymann D., Walton J.R., Jordan J.L., Lakatos S. and Yaniv A. (1975) Light and dark soils at the Apollo 16 landing site. *The Moon* 13, 81-110.
- Houck Karen (1982) Petrologic variations in Apollo 16 surface soils. *Proc. 13th Lunar Sci. Conf.* A197-A209.
- Imamura M., Nishiizumi K., Honda M., Finkel R.C., Arnold J.R. and Kohl C.P. (1974) Depth profiles of ⁵³Mn in lunar rocks and soils. *Proc. 5th Lunar Sci. Conf.* 2093-2104. 61221 61241 63321
- Jovanovic S. and Reed G.W. (1973) Volatile trace elements and the characterization of the Cayley formation and the primitive lunar crust. *Proc. 4th Lunar Sci. Conf.* 1313-1324.
- Kerridge J.F., Kaplan I.R. and Petrowski C. (1975) Nitrogen in the lunar regolith: Solar origin and effects. (abs) *LS VI*, 469-471.
- Kerridge J.F., Kaplan I.R. and Petrowski C. (1975) Evidence for meteoritic sulfur in lunar regolith. *Proc. 6th Lunar Sci. Conf.* 2151-2162.
- Korotev R.L. (1981) Compositional trends in Apollo 16 soils. *Proc. 12th Lunar Planet. Sci.* 577-605.
- Krahenbuhl U., Ganapathy R., Morgan J.W. and Anders E. (1973b) Volatile elements in Apollo 16 samples – Implications for highland volcanism and accretion history of the Moon. *Proc. 4th Lunar Sci. Conf.* 1325-1348.
- LSPET (1973) The Apollo 16 samples – petrographic and chemical description. *Science* 179, 23-34.
- Mark R.K., Cliff R.A., Lee-Hu C. and Weatherill G.W. (1973) Rb-Sr studies of lunar breccias and soils. *Proc. 4th Lunar Sci. Conf.* 1785-1796.
- Marvin U.B. (1972) Apollo 16 coarse fines (4-10 mm): Sample classification, description and inventory.
- Marvin U.B. and Mosie A.B. (1980) Apollo 16 soil catalog 61220: Classification and description of 1-4 mm fines. *JSC Curatorial Pub.* 53
- Marvin U.B. and Warren P.H. (1980) A pristine eucrite-like gabbro from Descartes and its exotic kindred. *Proc. 11th Lunar Planet. Sci. Conf.* 507-521.
- Meyer C. (1978) Ion microprobe analyses of aluminous lunar glasses: A test of the “rock type” hypothesis. *Proc. 9th Lunar Planet. Sci. Conf.* 1551-1570.
- Moore C.B., Lewis C.F. and Gibson E.K. (1973) Total carbon contents of Apollo 15 and 16 lunar samples. *Proc. 4th Lunar Sci. Conf.* 1613-1623.
- Morris R.V. (1978) The surface exposure (maturity) of lunar soils: Some concepts and Is/FeO compilation. *Proc. 9th Lunar Sci. Conf.* 2287-2298.
- Morris R.V., Score R., Dardano C. and Heiken G. (1983) *Handbook of Lunar Soils*. TWO volumes! *JSC* 19069.
- Nautiyal C.M., Padia J.T., Rao M.N. and Venkatesan T.R. (1981) Solar flare Neon: Clues from implanted noble gases in lunar soils and rocks. *Proc. 12th Lunar Planet. Sci. Conf.* 627-637.
- Nyquist L.E., Hubbard N.J., Gast P.W., Bansal B.M., Wiesmann H. and Jahn B.M. (1973) Rb-Sr systematics for

chemically defined Apollo 12 and 16 materials. Proc. 4th Lunar Sci. Conf. 1823-1846.

Rao M.N., Venkatesan T.R., Goswami J.N., Nautiyal C.M. and Padia J.T. (1979) Noble gas based solar flare exposure history of lunar rocks and soils. Proc. 10th Lunar Planet. Sci. Conf. 1547-1564.

Rhodes J.M., Adams J.B., Blanchard D.P., Charette M.P., Rodgers K.V., Jacobs J.W., Brannon J.C. and Haskin L.A. (1975) Chemistry of the agglutinate fractions in lunar soils. Proc. 6th Lunar Sci. Conf. 2291-2308.

Ridley W.I., Reid A.M., Warner J.L., Brown R.W., Gooley R. and Donaldson C. (1973) Glass compositions in Apollo 16 soils 60501 and 61221. Proc. 4th Lunar Sci. Conf. 309-321.

Rose H.J., Cuttitta F., Berman S., Carron M.K., Christian R.P., Dwornik E.J., Greenland L.P. and Ligon D.T. (1973) Compositional data for twenty-two Apollo 16 samples. Proc. 4th Lunar Sci. Conf. 1149-1158.

Ryder G. and Norman M.D. (1980) Catalog of Apollo 16 rocks. Part 1. JSC 16904

Sill G.T., Nagy B., Nagy L.A., Hamilton P.B., McEwan W.S. and Urey H.C. (1974) Carbon compounds in Apollo 17 lunar samples: Indications of cometary contribution to breccia 78155 ? (abs) LS V, 703-705.

Simoneit B.R., Christiansen P.C. and Burlingame A.L. (1973a) Volatile element chemistry of selected lunar, meteoritic and terrestrial samples. Proc. 4th Lunar Sci. Conf. 1635-1650.

Simoneit B.R., Christiansen P.C., Jackson R.F., Holland P.T. and Burlingame A.L. (1973b) Volatile element chemistry of selected lunar, meteoritic and terrestrial samples. (abs) LS IV, 679-681.

Sutton R.L. (1981) Documentation of Apollo 16 samples. Page 231. In USGS Prof. Paper 1048 (Ulrich et al. eds.)

Takeda H., Mori H., Ishii T. and Miyamoto M. (1981) Thermal and impact histories of pyroxenes in lunar eucrite-like gabbros and eucrites. Proc. 12th Lunar Planet. Sci. Conf. 1297-1313.

Taylor H.P. and Epstein S. (1973) O18/O16 and Si30/Si28 studies of some Apollo 15, 16 and 17 samples. Proc. 4th Lunar Sci. Conf. 1657-1679.

Walton J.R., Lakatos S. and Heymann D. (1973) Distribution of inert gases in fines from the Cayley-Descartes region. Proc. 4th Lunar Sci. Conf. 2079-2095.

Wszolek P.C., Simoneit B.R. and Burlingame A.L. (1973) Studies of magnetic fines and volatile-rich soils: Possible meteoritic and volcanic contributions to lunar carbon and light element chemistry. Proc. 4th Lunar Sci. Conf. 1693-1706.